


## 2011 PreAP Momentum 1

Variable	Units	Variable Name	Notes:
p (small)	kgm/s	momentum	How hard it is to stop something. Can be neg or 0.
I	kgm/s or Nsec	Impulse	Causes a change of p.


$p = mv$
$I = Ft$
$p_{\text{net}} = p_1 + p_2 \dots$

- \* A 35 kg object has  $-450$  kgm/s of momentum. Calculate its velocity.
- An object has 5000 kgm/s of momentum when it is moving 25 m/s. Calculate its mass.
- Which has more momentum? (*choose one for each*)
  - A car when going fast or slow?
  - A heavy or light object going 10 m/s?
- Which of the following has the most inertia?
  - \* A car when going fast or slow?
  - A heavy or light object going 10 m/s?
- Find the momentum of each of the following objects:




A. \_\_\_\_\_ B. \_\_\_\_\_ C. \_\_\_\_\_

- Which of the objects in #5 has the momentum with the greatest **magnitude** (*disregarding direction*)?
- Which of the objects in #5 has the most **inertia**?
- \* Find the net momentum (total) of all of the objects in #5 above (*find  $\Sigma p$* ).
- A 10 kg object is 5 m/s moving to the left while a 3 kg object is going 4 m/s to the right. (*Remember that left is negative.*)
  - Find the momentum of the 10 kg object (we'll call this momentum 1 or " $p_1$ "):
    - Find the momentum of the 3 kg object ( $p_2$ ):
    - Find the net momentum of both objects ( $\Sigma p$ ).
- \* A 25 kg object moving 3 m/s to the right while a 30 kg object is moving 4 m/s to the right (yes, same direction). Calculate  $p_{\text{net}}$ .
- A 2 kg object initially going 4 m/s to the right is later going 8 m/s. Find  $\Delta v$ . (*Remember that  $\Delta = \text{final} - \text{initial}$ .*)
- \* A 3 kg object going 6 m/s to the right ends up going 3 m/s to the left. Being careful of negatives and positives, find the change of momentum of the object.




**Lecture time:** In the last chapter Work caused a change of energy because the units for work are the same as for energy: joules.

It turns out that  $Ft$  (force times time) has the same units as momentum. Therefore: an impulse causes a change of momentum.



- 11) Slim Jim pushes on a 4 kg box for 3 seconds.
- A. Under the diagram, calculate the momentum before and after and the impulse Jim gave to the box.
- B. \* What does the impulse equal?



- 12) This time Slim Jim pushes on an object that was already moving.
- A. Under the diagram, calculate the momentum before and after and the impulse Jim gave to the box.
- B. What does the impulse equal?

So, this is our equation:  $\Sigma p_{\text{before}} \pm I = \Sigma p_{\text{after}}$ . Again, this is the same as in energy, where:  $\Sigma E_{\text{before}} \pm W = \Sigma E_{\text{after}}$ .

Q1: -12.9 m/s      Q4A: inertia is only about mass, so "same"

Q6: -30kgm/s (add 'em up).

Q8: 195 kgm/s

Q10: -27kgm/s =  $p_{\text{final}} - p_{\text{initial}}$

Q11B:  $I = p_{\text{final}}$