


Name: _____


Period: _____

Calculating net E, F, PE, and V


Key on back




1. Calculate the four electrostatic quantities (E, F, PE, and V) at a position 3 mm to the right of a $7\mu\text{C}$ charge. Be sure to give direction for vectors. Some quantities may be zero.



2. Calculate the four electrostatic quantities at a point 5 mm to the left of a $-5\mu\text{C}$ charge.




3. Now put the two previous problems together. Using the numbers you found in Q1 and 2, find the net electric field, net voltage, net force, and net energy at point P due to both charges. Again, some may be zero.



4. A $1.2\mu\text{C}$ charge is then brought to point P from infinity.
A. Again, using your previous numbers, calculate the four electrostatic quantities for this charge at point P.

B. How much work was done to move the charge to point P from infinity?



5. Now the negative charge is moved to the positive y-axis. Using the same individual numbers you calculated in Q2 and Q3, calculate the four quantities at point P.


6. A. Which way will the $1.2\mu\text{C}$ charge move when released?

B. If a negative charge was put at P, which way would it move?

Name: _____
 Period: _____

Calculating net E, F, PE, and V

remember to use a + test charge




$$\vec{E} = \frac{k(7 \times 10^{-6})}{(3 \times 10^{-3})^2} = 7 \times 10^9 \frac{N}{C} \text{ to right}$$

$$V = \frac{k(7 \times 10^{-6})}{3 \times 10^{-3}} \text{ or } E r = (7 \times 10^9) 3 \times 10^{-3} = 2.1 \times 10^7 \text{ J/C (no direction) (scalar)}$$

PE and $\vec{F} = 0$, since only 1 charge.

1. Calculate the four electrostatic quantities (E, F, PE, and V) at a position 3 mm to the right of a 7 μC charge. Be sure to give direction for vectors. Some quantities may be zero.




$$\vec{E} = \frac{k(5 \mu C)}{(5 \text{ mm})^2} = 1.8 \times 10^9 \text{ N/C to right}$$

$$V = E r = (1.8 \times 10^9)(5 \times 10^{-3}) = 9 \times 10^6 \text{ J/C (no direction)}$$

PE = $\vec{F} = 0$ (only 1 q)

neg. charges have neg. voltages

2. Calculate the four electrostatic quantities at a point 5 mm to the left of a -5 μC charge.




$$\vec{E}_2 + \vec{E}_5, \text{ since same dir, so } \vec{E}_{net} = (1.8 \times 10^9) + (7 \times 10^9) = 8.8 \times 10^9 \text{ N/C} \rightarrow$$

$$V_{net} = (-9 \times 10^6) + (2.1 \times 10^7) = 1.2 \times 10^7 \text{ J/C}$$

3. Now put the two previous problems together. Using the numbers you found in Q1 and 2, find the net electric field, net voltage, net force, and net energy at point P due to both charges. Again, some may be zero.

F_{net} and PE = 0, of course. Nothing at the position, so no F or PE.



4. A 1.2 μC charge is then brought to point P from infinity. A. Again, using your previous numbers, calculate the four electrostatic quantities for this charge at point P.


Just use net V and net E, which don't change
 E is in N/C, so $\vec{F} = q\vec{E}$, so $(1.2 \times 10^{-6}) 8.8 \times 10^9 \frac{N}{C} = 1.056 \times 10^4 \text{ N to R} \rightarrow$
 (since +s go dir. of \vec{E})
 PE is in J, so PE = qV, so $(1.2 \times 10^{-6} \text{ C}) (1.2 \times 10^7 \text{ J/C}) = 14.4 \text{ J}$

B. How much work was done to move the charge to point P from infinity?

$$PE = KE = W = 14.4 \text{ J}$$

5. Now the negative charge is moved to the positive y-axis. Using the same individual numbers you calculated in Q2 and Q3, calculate the four quantities at point P.

Easiest to calculate V_{net} and E_{net} , then just multiply by the q at point P. That way you don't have to do pyth and inverse tan twice.



V_{net} = same, since it is a scalar and dir. doesn't matter
 $V_{net} = 1.2 \times 10^7 \text{ J/C}$

$$\vec{E} = 1.8 \times 10^9 \frac{N}{C} \uparrow$$

$$V = -9 \times 10^6 \text{ J/C}$$


$$(\vec{E}_{net})^2 = (1.8 \times 10^9)^2 + (7 \times 10^9)^2 \text{ (can do without exponents. All are } \times 10^9)$$

$$\vec{E}_{net} = 7.23 \times 10^9 \text{ N/C}$$

$$\text{and } \vec{F} = q\vec{E} = [(1.2 \times 10^{-6}) (7.23 \times 10^9)] = 8.68 \times 10^3 \text{ N at } 14.4^\circ$$

$$\theta = \tan^{-1}(y/x) = 14.4^\circ$$

$$[N] = [C][N/C]$$



6. A. Which way will the 1.2 μC charge move when released?
 at 14.4°, same way \vec{E} points.

B. If a negative charge was put at P, which way would it move?

at 14.4 + 180 = 194.4°, negs move opp. dir. of electric field

since PE = qV and is also a scalar, it stays same = 14.4 J